This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problems Mailbox.

FIG. 1



1

FIG. 2






FIG. 3




I lag phase

- II rapid wound contraction phase
- III plateau phase



Single and double asterisks signify that the STZ-Diabetic group (n=5) differed from the normal-untreated group (n=10) with a P< 0.05 and P< 0.005 by Mann Whitney test (Ttest), respectively.



Days after wound induction

Single and double asterisks signify that the STZ-Diabetic group (n=5) differed from the normal-untreated group (n=10) with a P< 0.05 and P< 0.005 by Mann Whitney test (T-test), respectively.

FIG. 5






FIG. 6A



n=5, Kruskal-wallis test p>0.05

FIG. 6B



n=5, Kruskal-wallis test p>0.05

- Normal, untreated
- Diabetic, treated with 676 after DM induction
- Normal, treated with 676 after wound induction







FIG. 8



FIG






FIG. 10A



mean blood gluc level, mg/dL 355

FIG. 10B



mean blood gluc level, mg/dL 381